


COVID-19 VACCINATION SCHEDULE FOR MODERATELY TO SEVERELY IMMUNOCOMPROMISED PATIENTS


Includes blood cancer patients and most survivors*

If you are moderately to severely immunocompromised, you are at increased risk of severe COVID-19 illness and death. Additionally, your immune response may not be as strong as in people who are not immunocompromised, so additional vaccine doses are needed. Even after receiving all recommended vaccines, LLS encourages blood cancer patients to take additional precautions, like wearing a mask and social distancing.

Pfizer-BioNTech



Moderna



Boosters may be “mixed and matched.”

Moderna boosters are a half-dose; Pfizer-BioNTech boosters are a regular dose.

Revaccination may be considered for patients who received one or more doses of COVID-19 vaccine while they were being treated with B-cell depleting therapies like rituximab or ocrelizumab. A patient’s own oncologist is the best person to determine whether and when revaccination should happen.

Moderna and Pfizer/BioNTech vaccines are strongly preferred over the Johnson & Johnson COVID-19 vaccine. **For information on doses and timing of the J&J vaccine**, please visit CDC at <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/different-vaccines/janssen.html>



Get vaccinated as recommended even if you had COVID-19



Keep practicing social distancing and masking



*Talk to your healthcare team if unsure of your risk