

# COVID-19 Vaccination Schedule

## for Moderately to Severely Immunocompromised Patients


Includes blood cancer patients and most survivors

If you are moderately to severely immunocompromised, you are at increased risk of severe COVID-19 illness and death. Additionally, your immune response to vaccination may not be as strong as in people who are not immunocompromised, so additional vaccine doses are needed. Even after receiving all recommended vaccines, LLS encourages blood cancer patients to take additional precautions, like wearing a mask and social distancing.


The charts below include information about the number and timing of COVID-19 vaccine doses for everyone 6 months of age and older. Individuals should consult with their healthcare provider about any changes to the number or timing of their doses due to their cancer diagnosis or its treatment, or receipt of monoclonal antibodies (you can find more information in our [COVID FAQs](#)).

### Moderna (Spikevax)

#### Ages 6 months – 17 years




#### Ages 18 years and older



**NOTE:** Regardless of health or immune status, everyone who is eligible should receive only one COVID-19 bivalent booster dose. Repeat dosing with bivalent boosters is not recommended at this time. LLS monitors new scientific information and public health recommendations closely and will update this information as needed.


## Pfizer-BioNTech (Comirnaty)

### Ages 6 months – 4 years




**NOTE ABOUT BIVALENT BOOSTER:** The first two primary doses should be with the monovalent vaccine. The third primary dose should be with the bivalent vaccine. Children in this age group who received three monovalent vaccines for their primary series should receive a bivalent booster dose at least two months after completing their primary series.

### Ages 5 – 11 years



**NOTE:** Regardless of health or immune status, everyone who is eligible should receive only one COVID-19 bivalent booster dose. Repeat dosing with bivalent boosters is not recommended at this time. LLS monitors new scientific information and public health recommendations closely and will update this information as needed.

### Ages 12 years and older



## Novavax

Ages 12 years and older



**NOTE:** Regardless of health or immune status, everyone who is eligible should receive only one COVID-19 bivalent booster dose. Repeat dosing with bivalent boosters is not recommended at this time. LLS monitors new scientific information and public health recommendations closely and will update this information as needed.



Get vaccinated as recommended even if you had COVID-19



Keep practicing social distancing and masking



Talk to your healthcare team if unsure of your risk